



## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.


Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

The intersections of the opposite sides are collinear, by the theorem of Pappus, that is, if a hexagon  $AB'CA'BC'$  has its vertices of odd order on one straight line, and its vertices of even order on another straight line, then the three pairs of opposite sides,  $AB'$  and  $A'B$ ,  $B'C$  and  $BC'$ ,  $CA'$  and  $C'A$ , meet in three points lying on another straight line.



But the intersections of the opposite sides of this hexagon are the intersections of the diagonals of the three quadrilaterals.

Hence, the intersections of the diagonals of any three quadrilaterals, two of which are formed by cutting the other one by a straight line, are collinear.

Also solved by ANNA MULLIKIN.

A solution of this problem appeared in the January issue of the MONTHLY, but we publish this solution as it presents an entirely different method of attack. EDITORS.

**452. Proposed by NATHAN ALTSHILLER, University of Washington.**

Through a given point a secant is drawn that meets three given concurrent lines in the points  $A, B, C$ , respectively. Determine the position of the secant by the condition  $AB/BC = K$ ,  $K$  being given.

**SOLUTION BY MRS. ELIZABETH B. DAVIS, U. S. Naval Observatory.**

Let  $OA', OB'$  and  $OC'$  be three given concurrent lines, and  $P$  a given point. Let it be required to draw through  $P$  a secant meeting  $OA', OB'$  and  $OC'$  respectively in points  $A, B$ , and  $C$ , such that  $AB/BC = K$ ,  $K$  being given.

Join  $P$  and  $O$ , and through  $P$  draw any transversal  $R'P$ , meeting the four lines of the pencil  $O - A'B'C'P$  in  $D, E, F$ , and  $P$ , respectively.

On  $OP$  take  $H$  and  $G$  so that

$$GP : HP = DE : EF. \quad (1)$$

Also, on  $OP$  take  $M$ , so that

$$MP : HP = K. \quad (2)$$

Join  $DG$ , and through  $M$  draw  $RM$  parallel to  $DG$ , meeting  $R'P$  in  $R$ .

Draw  $RA$  parallel to  $OC'$ , meeting  $OA'$  in  $A$ . Join  $AP$ , then  $AP$  is the transversal required. For, dividing (1) by (2), we have

$$GP : MP = \frac{DE}{EF} : K.$$

Since,  $\triangle$ 's  $PDG$  and  $PRM$  are similar,

$$GP : MP = DP : RP.$$

Hence


$$DP : RP = \frac{DE}{EF} : K. \quad (3)$$

Dividing the first ratio of (3) by  $FP$ ,

$$\frac{DP}{FP} : \frac{RP}{FP} = \frac{DE}{EF} : K. \quad (4)$$

Since  $\triangle$ 's  $RAP$  and  $FCP$  are similar,

$$\frac{RP}{FP} = \frac{AP}{CP}.$$



Substituting this value of  $RP/FP$  in (4), we have

$$\frac{DP}{FP} : \frac{AP}{CP} = \frac{DE}{EF} : K,$$

or

$$\frac{DP}{FP} : \frac{DF}{EF} = \frac{AP}{CP} : K. \quad (5)$$

But the anharmonic ratio of all transversals cutting any given pencil of four lines, as  $O-A'B'C'P$ , is a constant. Hence,

$$\frac{DP}{FP} : \frac{DE}{EF} = \frac{AP}{CP} : \frac{AB}{BC}. \quad (6)$$

From (5) and (6),  $AB/BC = K$ . Hence  $AP$  is the transversal required.

Also solved by the PROPOSER.

**453. Proposed by CLIFFORD N. MILLS, Brookings, S. D.**

Prove geometrically the formulæ for  $\sin 2\beta$ ,  $\cos 2\beta$ ,  $\sin 3\beta$ ,  $\cos 3\beta$ .

SOLUTION BY THE PROPOSER.

$$(1) \sin 2\beta = 2 \sin \beta \cos \beta.$$

Inscribe in a circle any triangle,  $ABC$ , with the angle at  $B$  equal to  $2\beta$  (Fig. 1). Draw  $AM$  through the center of the circle;  $BK$ , the bisector of the angle  $B$ ;  $OK$ , the radius of the circle;  $AK$ ,  $KM$ , and  $KC$ ,  $K$  being on the circumference of the circle.

Then

$$\sin 2\beta = \frac{AC}{2r}.$$

$\triangle AKC$  and  $KOM$  are similar isosceles triangles. Hence

$$\frac{AC}{AK} = \frac{KM}{r} \quad \text{or} \quad AC = \frac{AK \cdot KM}{r}.$$

Hence,

$$\frac{AC}{2r} = \frac{AK \cdot KM}{2r^2} = \frac{2AK \cdot KM}{4r^2}.$$

But

$$\frac{AK}{2r} = \sin \beta \quad \text{and} \quad \frac{KM}{2r} = \cos \beta.$$

Hence  $\sin 2\beta = 2 \sin \beta \cos \beta$ .

$$(2) \cos 2\beta = \cos^2 \beta - \sin^2 \beta.$$

Using the same Fig. 1, we get

$$\cos 2\beta = \frac{MC}{2r} = \sqrt{\frac{4r^2 - AC^2}{4r^2}} = \sqrt{1 - \frac{AC^2}{4r^2}}.$$

But

$$AC = (AK \cdot KM)/r.$$

Hence,

$$\cos 2\beta = \sqrt{1 - \frac{4AK^2 \cdot KM^2}{16r^4}}.$$

Substituting the values of  $AK$  and  $KM$ , as previously found, we have

$$\cos 2\beta = \sqrt{1 - 4 \sin^2 \beta \cos^2 \beta} = \sqrt{(\sin^2 \beta + \cos^2 \beta)^2 - 4 \sin^2 \beta \cos^2 \beta} = \cos^2 \beta - \sin^2 \beta.$$

$$(3) \sin 3\beta = 3 \sin \beta - 4 \sin^3 \beta.$$




Fig. 1.